

Complete Genome Sequence of *Serratia marcescens* SmUNAM836, a Nonpigmented Multidrug-Resistant Strain Isolated from a Mexican Patient with Obstructive Pulmonary Disease

Luisa Sandner-Miranda,^a Pablo Vinuesa,^b Gloria Soberón-Chávez,^c Rosario Morales-Espinosa^a

Facultad de Medicina,^a Centro de Ciencias Genómicas,^b and Instituto de Investigaciones Biomédicas,^c Universidad Nacional Autónoma de México, Mexico City, Mexico

L.S.-M. and P.V. contributed equally to this work.

***Serratia marcescens* SmUNAM836 is a multidrug-resistant clinical strain isolated in Mexico City from a patient with chronic obstructive pulmonary disease. Its complete genome sequence was determined using PacBio RS II SMRT technology, consisting of a 5.2-Mb chromosome and a 26.3-kb plasmid, encoding multiple resistance determinants and virulence factors.**

Received 12 October 2015 Accepted 16 October 2015 Published 21 January 2016

Citation Sandner-Miranda L, Vinuesa P, Soberón-Chávez G, Morales-Espinosa R. 2016. Complete genome sequence of *Serratia marcescens* SmUNAM836, a nonpigmented multidrug-resistant strain isolated from a Mexican patient with obstructive pulmonary disease. *Genome Announc* 4(1):e01417-15. doi: 10.1128/genomeA.01417-15.

Copyright © 2016 Sandner-Miranda et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Rosario Morales-Espinosa, marosari@unam.mx.

Serratia marcescens is emerging as an opportunistic nosocomial pathogen causing urinary tract, bloodstream, skin, and respiratory infections (1). Several virulence factors have been identified, such as hemolysins, proteases, and siderophores (2). Many clinical isolates of this species display intrinsic or acquired resistance to multiple antibiotic families (1).

Here we present the complete genome sequence of *Serratia marcescens* strain SmUNAM836, a nonpigmented, multidrug-resistant bacterium isolated from the bronchial aspiration of a hospitalized patient with chronic obstructive pulmonary disease in Mexico City, as detailed in BioProject accession no. PRJNA284857. Strain SmUNAM836 was resistant to diverse β -lactams, including penicillins, but it is susceptible to cephalosporins and carbapenems. It is resistant to aminoglycosides (to-bramycin), fluoroquinolones (ciprofloxacin), and polymixin B.

The genomic DNA of the strain was purified with the DNeasy blood and tissue kit (Qiagen) and sent to the Yale Center for Genome Analysis (YCGA) for PacBio RS II single-molecule real-time (SMRT) sequencing. A standard library of 10-kb fragments was prepared and sequenced on three SMRT cells with the P2-C4 chemistry. The continuous long reads were assembled using the HGAP/Quiver protocol in SMRT portal v.2.3.0.140936.p4 (3), resulting in an assembly with 2 contigs. They were circularized by trimming the terminal repeats with Minimus2 (4) and subjected to two consecutive rounds of read remapping with the RS_Resequencing.1 module for sequence polishing. The final assembly had a mean coverage of $\sim 172\times$ and consists of a chromosome (5,207,023 bp), and a plasmid (26,346 bp), with a mean G+C content of 59.71%. Gene calling and annotation were performed with a modified version of Prokka (5). A total of 4,961 genes were annotated: 4788 CDSs, 95 tRNAs, 22 rRNAs, and 1 tmRNA. Additionally, 45 ncRNAs, 3 riboswitches, and 502 signal peptides were also identified, but no clustered regularly interspaced short palindromic repeat (CRISPR) arrays. The annotation was manually curated and enriched by adding 6 prophage predictions made

by the PHAST server (6) and 13 genomic islands identified by IslandViewer3 (7).

The genome was screened for the presence of antibiotic resistance genes, efflux pumps, and potential virulence factors using blastp (8) searches against locally maintained versions of the CARD (9), ResFinder (10), and VirulenceFinder (11) databases. Eight resistance-nodulation-division (RND), two ATP-binding cassette (ABC), and one major-facilitator-superfamily (MFS) type efflux pumps were identified, along with the chromosomally encoded BlaKPC-2 (class B) and AmpC-like (class C) beta-lactamases, the aminoglycoside acetyltransferase Aac(6')-Ic, and the aminoglycoside phosphotransferase StrA, an QnrB31-like fluoroquinolone resistance protein. No class 1 or class 2 integrons or ISCR elements were found.

The plasmid had a low G+C content (43.52%) and displayed $\sim 97\%$ sequence identity with diverse *Enterobacter cloacae* plasmids like pKPC-f91. It harbors the transposon Tn3 encoding 3 proteins: a BlaTEM-1D (class A) beta-lactamase and the Tn3 transposase and resolvase. It encodes the genes for the *ccdAB* toxin-antitoxin system.

Potential virulence factors detected are long polar fimbriae (LpfA), hemolysin D (HlyD), and the ClpP protease, the Iron enterobactin siderophore receptor, hemolytic phospholipase C (PlcH), and plasmid-encoded catalase peroxidase (KatP) and enterotoxin (SenB).

Nucleotide sequence accession numbers. The complete genome sequence of *Serratia marcescens* SmUNAM836 is available in GenBank under accession numbers CP012685 (chromosome) and CP012686 (plasmid), BioProject accession number PRJNA284857, and BioSample accession number SAMN03733572.

ACKNOWLEDGMENTS

We thank Gabriela Delgado for her academic support and Biol. José Luis Méndez for performing the antibiotic susceptibility essays. Guilin Wang from the Yale Center for Genome Analysis, Yale University, is acknowl-

edged for skillful PacBio sequencing. Alfredo Hernández Álvarez, from the Centro de Ciencias Genómicas, UNAM, is gratefully acknowledged for support with server system administration. This paper is part of the doctoral research carried out by L. Sandner Miranda and constitutes partial fulfillment of the “Programa de Doctorado en Ciencias Biomédicas, Universidad Nacional Autónoma de México (UNAM).”

This work was supported by CONACYT-México (100343 to R.M.E. and 179133 to P.V.) and DGAPA/PAPIIT-UNAM (IN212513 to R.M.E. and IN211814 to P.V.). The funders had no role in study design, data collection and interpretation, or decision to submit the work for publication.

FUNDING INFORMATION

Dirección General Asuntos del Personal Académico, Universidad Nacional Autónoma de México (DGAPA, UNAM) provided funding to Rosario Morales-Espinosa under grant number IN212513. Dirección General Asuntos del Personal Académico, Universidad Nacional Autónoma de México (DGAPA, UNAM) provided funding to Pablo Vinuesa under grant number IN211814. Consejo Nacional de Ciencia y Tecnología (CONACYT) provided funding to Rosario Morales-Espinosa under grant number 100343. Consejo Nacional de Ciencia y Tecnología (CONACYT) provided funding to Pablo Vinuesa under grant number 179133.

REFERENCES

- Iguchi A, Nagaya Y, Pradel E, Ooka T, Ogura Y, Katsura K, Kurokawa K, Oshima K, Hattori M, Parkhill J, Sebaihia M, Coulthurst SJ, Gotoh N, Thomson NR, Ewbank JJ, Hayashi T. 2014. Genome evolution and plasticity of *Serratia marcescens*, an important multidrug-resistant nosocomial pathogen. *Genome Biol Evol* 6:2096–2110.
- Shimuta K, Ohnishi M, Iyoda S, Gotoh N, Koizumi N, Watanabe H. 2009. The hemolytic and cytolytic activities of *Serratia marcescens* phospholipase A (PhlA) depend on lysophospholipid production by PhlA. *BMC Microbiol* 9:261.
- Chin C, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
- Treangen TJ, Sommer DD, Angly FE, Koren S, Pop M. 2012. Next generation sequence assembly with AMOS. *Curr Protoc Bioinformatics Chapter 11:Unit 11:8*. <http://dx.doi.org/10.1002/0471250953.bi1108833>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
- Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res* 39:W347–W352. <http://dx.doi.org/10.1093/nar/gkr485>.
- Dhillon BK, Laird MR, Shay JA, Winsor GL, Lo R, Nizam F, Pereira SK, Waglechner N, McArthur AG, Langille MGI, Brinkman FSL. 2015. IslandViewer 3: More flexible, interactive genomic island discovery, visualization and analysis. *Nucleic Acids Res* 43(Web Server Issue):W104–W108. <http://dx.doi.org/10.1093/nar/gkv401>.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <http://dx.doi.org/10.1186/1471-2105-10-421>.
- McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJ, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang YM, Yu T, Wright GD. 2013. The comprehensive antibiotic resistance database. *Antimicrob Agents Chemother* 57:3348–3357. <http://dx.doi.org/10.1128/AAC.00419-13>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.
- Kleinheinz KA, Joensen KG, Larsen MV. 2014. Applying the ResFinder and VirulenceFinder web-services for easy identification of acquired antibiotic resistance and virulence genes in bacteriophage and prophage nucleotide sequences. *Bacteriophage* 4:e27943. <http://dx.doi.org/10.4161/bact.27943>.